


---

# Insect Sex Pheromones

---

Martin Jacobson

---



---

Academic Press

---

## **INSECT SEX PHEROMONES**

This page intentionally left blank

# INSECT SEX PHEROMONES

MARTIN JACOBSON

*Agricultural Environmental Quality Institute  
Agricultural Research Service  
U.S. Department of Agriculture  
Beltsville, Maryland*



1972

ACADEMIC PRESS New York and London

**COPYRIGHT © 1972, BY ACADEMIC PRESS, INC.**

**ALL RIGHTS RESERVED.**

**NO PART OF THIS PUBLICATION MAY BE REPRODUCED OR TRANSMITTED IN ANY FORM OR BY ANY MEANS, ELECTRONIC OR MECHANICAL, INCLUDING PHOTOCOPY, RECORDING, OR ANY INFORMATION STORAGE AND RETRIEVAL SYSTEM, WITHOUT PERMISSION IN WRITING FROM THE PUBLISHER.**

**ACADEMIC PRESS, INC.**

**111 Fifth Avenue, New York, New York 10003**

*United Kingdom Edition published by*  
**ACADEMIC PRESS, INC. (LONDON) LTD.**  
**24/28 Oval Road, London NW1**

**LIBRARY OF CONGRESS CATALOG CARD NUMBER: 72-77360**

**PRINTED IN THE UNITED STATES OF AMERICA**

*To*

DEBRA GAIL *and* LINDA HOPE

This page intentionally left blank

## **CONTENTS**

*Preface*    *xi*

### **Chapter I Introduction**

Text    *1*

### **Chapter II Occurrence of Sex Pheromones in the Female**

Acarina	<i>5</i>
Orthoptera	<i>6</i>
Hemiptera	<i>7</i>
Homoptera	<i>8</i>
Diptera	<i>9</i>
Isoptera	<i>12</i>
Neuroptera	<i>12</i>
Siphonaptera	<i>12</i>
Coleoptera	<i>12</i>
Hymenoptera	<i>19</i>
Lepidoptera	<i>27</i>

### **Chapter III Sex Pheromones Produced by Males**

Acarina	<i>55</i>
Orthoptera	<i>55</i>
Hemiptera	<i>56</i>
Trichoptera	<i>57</i>

Coleoptera	57
Hymenoptera	60
Diptera	61
Lepidoptera	63
Mecoptera	69

#### **Chapter IV Pheromones Produced by One Sex That Lure Both Sexes (Assembling or Aggregating Scents)**

Orthoptera	71
Hemiptera	72
Coleoptera	72

#### **Chapter V Anatomy and Physiology of the Production Glands**

In Females	79
Orthoptera	79
Lepidoptera	80
Coleoptera	90
Hemiptera (Homoptera)	92
Hymenoptera	92
Diptera	92
In Males	93
Orthoptera	93
Lepidoptera	93
Coleoptera	98
Hemiptera (Heteroptera)	99
Diptera	99
Mecoptera	100

#### **Chapter VI The Mechanism of Attractant Perception**

Text	101
------	-----

#### **Chapter VII Responses to Sex Pheromones**

Behavioral Responses	122
Acarina	122
Orthoptera	122

Hemiptera	126
Diptera	128
Coleoptera	129
Hymenoptera	133
Lepidoptera	133

Electrophysiological Responses	144
--------------------------------	-----

### **Chapter VIII Influence of Age of the Insect on Production of and Response to Sex Pheromones**

Orthoptera	149
Hemiptera (Heteroptera)	149
Hemiptera (Homoptera)	150
Lepidoptera	150
Coleoptera	153
Hymenoptera	154
Neuroptera	154
Diptera	154

### **Chapter IX Influence of Time of Day on Sex Pheromone Production and Mating**

Orthoptera	157
Lepidoptera	158
Coleoptera	161
Hymenoptera	161
Diptera	161

### **Chapter X Collection, Isolation, and Identification of Sex Pheromones**

Collection	163
Isolation	164
Identification	164
Acarina	165
Orthoptera	165
Hemiptera	168
Lepidoptera	170
Coleoptera	193
Hymenoptera	205
Diptera	207

**Chapter XI Synthesis of the Sex Pheromones**

Text 209

**Chapter XII Sex Attractants in Insect Survey**

Text 243

**Chapter XIII Sex Pheromones in Insect Control**

Text 277

**Bibliography** 293*Subject Index* 361

## **PREFACE**

The first edition of this reference work, "Insect Sex Attractants," was published in 1965 by Wiley (Interscience). To date it is still the only book of its kind on the subject. However, the surge of interest in the insect sex pheromones as evidenced by the tremendous progress and by the phenomenal additional publications since 1965 has made it imperative to bring the entire subject up-to-date in this revised and expanded version.

In my earlier Preface I stated, "It is my sincere hope that this book will provide an incentive for greater discoveries in this fascinating field." An unprecedented increase in the number of species shown to produce sex pheromones (more than double the number known in 1965), in the number of species whose sex pheromones have been identified (37 now as compared to 3 in 1965), and the extensive literature in the field (about 1400 references as compared to 425 in 1965) indicate that this wish has been fulfilled.

The discovery that many sex pheromones are sexually excitatory rather than attractive has prompted me to substitute the more accurate and encompassing term "pheromones" for the term "attractants" in the title of this edition. Not only have all of the original chapters been extensively enlarged and revised, but a new chapter on test methods and responses has been added. This expanded review of the world literature should be extremely valuable to research and economic entomologists, insect physiologists, chemists, and ecologists.

Photographs and other illustrations are used with the kind permission of the copyright owners, with the source of each indicated. All listed references have been consulted directly. References which have appeared or were noted after the manuscript was prepared have been assigned

supplementary numbers, enabling me to include literature which appeared up to a few months prior to publication.

I acknowledge a debt of gratitude to my wife, Nettie, for painstakingly typing the bulk of the manuscript. A special note of thanks is due a number of experts on insect taxonomy at the National Museum of the Smithsonian Institution, Washington, D.C., for checking many of the insect names.

Martin Jacobson

## CHAPTER I

### INTRODUCTION

Insects have managed to persist in hostile surroundings because they have developed extraordinary adaptations or abilities, one of which is a highly specialized sense of smell. Because many insects depend on their sense of smell for survival, they can frequently be attracted to a trap by a chemical for detection purposes, to a toxicant that destroys them, or to a substance that makes them incapable of fertile mating (599).

Attractants may be classified as sex, food, or oviposition lures. The type of lure is inferred or deduced from insect behavior, and assignment is frequently uncertain. A chemical is probably a sex attractant if it brings to it an insect, which then assumes a mating position or attempts to mate with the chemical or with an object on which the chemical has been placed. This definition excludes a number of substances, mainly food attractants, which lure only one sex but do not elicit a sexual response. Sex attractants, usually released by a female to lure a male, are important, if not essential, links in the process by which the sexes locate each other for mating. Although odors released by female insects are usually for the purpose of attracting males from a distance, they may also serve to sexually excite the male before copulation and to evoke a courtship response. Sexual odors released by males are primarily for the purpose of sexually exciting the female, making her more receptive to the male's advances (aphrodisiacs). However, species

are known in which the males produce distance attractants for the female.

These chemical messengers are also called "assembling scents" (663) and "sex pheromones," from the Greek *pherein* (to carry) and *horman* (to excite, stimulate) (647). The term pheromone has been attacked by Kirschenblatt (676) as being etymologically incorrect, as it gives no clue to its exact derivation. Kirschenblatt (674, 675) previously proposed the term "telergones," from the Greek *tele* (afar) and *ergon* (action), to designate all biologically active substances secreted by animals into their environment which influence other organisms ("these substances are products of external secretion and differ principally by their importance from hormones, which display their physiological action within the organism producing them") (674). Micklem (787) cited the same objection to the term "pheromone" and suggested that it be changed to "pherormone." In replying to Micklem, Karlson and Lüscher (649) gave a new etymological explanation for their term, stating that the ending "mone" is regarded as a proper suffix used in such scientific terms as "hormones," "gamones," and "termones." "Pheromone" is now commonly used and widely accepted to include those substances secreted by an animal to influence the behavior of other animals of the same species (1268). Brown *et al.* (213) have recently proposed the term "allomone" to include those chemical substances produced by an organism which evoke a behavioral or physiological reaction in an organism of another species, and the term "kairomone" as a transspecific chemical messenger of benefit to the recipient rather than to the producer.

The use of sex pheromones by organisms other than insects has been conclusively demonstrated in algae (805, 878), nematodes (484, 485), spiders (532), crustaceans (77, 323, 679, 1011), fishes (1187), and mammals such as dogs (354), cattle (354), deer (810), mice (158, 1129), hamsters (814), and primates (321, 784-786, 1237). Indeed, it is possible that sex pheromones play a part in the courtship and reproduction of humans as well (66, 306, 1177a). Excellent reviews of this subject are those by Whitten (1263), Michael and Keverne (784), Bruce (214, 215), and Ralls (906).

General reviews on various aspects of the subject of insect pheromones may be found in references 112, 144-146, 151-154, 229, 244, 247, 305, 375, 375a, 422, 468, 479, 501, 521, 525, 529, 530, 545, 550, 574, 587, 594, 596, 599, 601, 636a, 639, 644, 669, 695, 700, 707, 713, 789, 807, 907, 932, 978, 979, 1092, 1156, 1188, 1254, 1268, 1272, and 1292. Brief reviews of insect sex pheromones are found in references 64, 78, 150, 192, 249, 316, 544, 590, 646, 650, 699, 738, 756, 797, 798, 811, 817, and 850. The assembling of various moths and butterflies has been reviewed by Poulton (883).

Reviews in references 37, 122, 128, 230, 592, 645, 703, 716, 812, 831a, 900 (140 references), 916, 1112, and 1131 deal mainly with the chemistry

of insect sex pheromones; reference *1079* deals with their specificity (or lack of it), and references *249*, *254*, *440*, *449*, and *1269* are devoted to honeybee pheromones. Sex pheromones among the Lepidoptera are reviewed mainly in references *540*, *606*, and *757*, and those among the Coleoptera are discussed in references *78*, *223*, *774*, *997*, *1100*, *1133a*, *1206*, and *1286*.

This page intentionally left blank

## CHAPTER II

### OCCURRENCE OF SEX PHEROMONES IN THE FEMALE

As long ago as 1837, von Siebold (1097) recognized that a pair of appendages, sometimes colored, opening into the vagina of the females of some insect species may act as an attractant for males. He surmised that the odor emitted by a female insect probably functions to entice the male, while that emitted by a male may be used as a stimulus in copulation (aphrodisiac).

#### ACARINA

*Amblyomma americanum* (L.), lone star tick

*Amblyomma maculatum* Koch, Gulf Coast tick

*Dermacentor variabilis* (Say), American dog tick

Females of these species produce a pheromone that attracts males of the respective species. Males respond only after reaching a state of maturity initiated by feeding (138).

*Panonychus ulmi* (Koch), European red mite

Males tend to aggregate around quiescent female deutonymphs to await the latter's eclosion (899).

*Tetranychus urticae* Koch, two-spotted spider mite

Males are attracted strongly to quiescent deutonymphs, remaining until emergence of the adult female, when mating occurs. They are also attracted to ether extracts of the deutonymphs (307, 308).

## ORTHOPTERA

*Blaberus craniifer* (Burmeister), giant death's head roach

Virgin females produce a volatile sex pheromone which attracts males and elicits antennal waving, alertness, and locomotion toward the females (107). It is also interspecifically effective in eliciting male courtship behavior in *Blaberus giganteus* and *Byrsotria fumigata*.

*Blaberus giganteus* L.

Virgin females may produce a volatile attractant for males, releasing courtship behavior (107).

*Blatta orientalis* L., oriental cockroach

Females appear to have a nonvolatile sex pheromone present on their body surface which facilitates sex recognition and releases male courtship behavior. The wing-raising display is less complex and variable than that of the male *Periplaneta americana* (114).

*Blattella germanica* L., German cockroach

Virgin females produce a substance attractive only to males. It is unattractive to males of *Blatta orientalis* and *Shelfordella tartara* (1237).

*Byrsotria fumigata* (Guérin)

Virgin females produce a volatile sex attractant which enables males to perceive them at a considerable distance and elicits alertness, antennal waving, and wing "pumping" (106, 107, 110, 981). A number of gynandromorphs are found to produce the female sex pheromone (116).

*Leucophaea maderae* (F.), Madeira cockroach

Smyth (1117) claims to have collected a volatile material from females that increases the incidence of courtship by males.

*Mantis religiosa* (L.), praying mantis

Caged virgin females can lure large numbers of males from a distance of up to 100 m between 8:30 AM and 1:00 PM (658).

*Nauphoeta cinerea* (Olivier), cockroach

The sex pheromone, if it exists, appears to be a nonvolatile substance on the surface of the female. Male display (raising of the wings) is readily evoked, even by unreceptive females (107).

*Periplaneta americana* (L.), American cockroach

Females emit an odorous attractant for the male. The substance adheres to paper or other materials with which the females come in contact (985). Virgin females, as well as filter papers exposed to them, cause male alertness, antennal movement, searching locomotion, and vigorous wing flutter (107, 985).

The wing-raising display, which is released much more readily in groups of males than in single males, is used as the single criterion of response in a bioassay method developed by Wharton *et al.* (1258, 1259). However,

Jacobson and Beroza (600) have shown that a number of organic compounds, including several that are repellent to males (such as amyl acetate), will elicit wing-raising, thus making it mandatory that an accurate bioassay show a combination of intense excitement, wing-raising, and attempts to copulate with one another. The mating urge is so powerful in this insect that males starved for 4 weeks in the laboratory and then given their choice of the female sex pheromone or food, invariably responded to the sex pheromone until they were near death (764).

The sex attractant is produced principally by virgin females and sporadically by mated females. Nymphs ordinarily do not produce the attractant, and a newly emerged female produces very little of the substance at first. During this nonproductive phase, the female does not attract the male, and mating does not occur. Maximum production is attained by the second week after eclosion. Carbon dioxide anesthetization and manual manipulation reduce the production of attractant somewhat (1260). Attractant synthesis is drastically depressed within 18 hours after copulation, which accounts for the sporadic production of the substance by mated females.

The female sex pheromone is effective in releasing courting behavior in males of other *Periplaneta* species and in males of *Blatta orientalis*, but not in *Eurycotis floridana*, *Leucophaea maderae*, or *Nauphoeta cinerea* (107). The sex pheromones of other species of *Periplaneta* also appear to be inter-specifically effective within the genus (114, 980).

*Periplaneta australasiae* (Fabr.), Australian cockroach

*Periplaneta brunnea* (Burmeister)

Females of these 2 species produce pheromones that attract and sexually excite males. The same behavior is elicited in males by exposure to filter papers over which females have crawled (107, 114).

*Periplaneta fuliginosa* (Serville)

Virgin females apparently produce a chemical substance which acts as a releaser of courtship behavior in males, but this has not yet been proved (107, 114).

## HEMIPTERA

*Dysdercus cingulatus* Fabr., red cotton bug

A cardboard box (with pinhole perforations) containing virgin females was placed 15 inches from a group of males. The males were quickly attracted to the box, and vibrated their wings and raised their antennae. A corresponding box without females did not attract males, but a box in which a female had been confined 5 days earlier acted as an attractant (837).