

Ernst Paunzen

The Δa photometric system



Cuvillier Verlag Göttingen
Internationaler wissenschaftlicher Fachverlag



The Δa photometric system





The Δa photometric system

Ernst Paunzen

Department of Theoretical Physics and Astrophysics
Masaryk University
Kotlářská 2
CZ - 611 37 Brno
Czech Republic





Bibliographical information held by the German National Library

The German National Library has listed this book in the Deutsche Nationalbibliografie (German national bibliography); detailed bibliographic information is available online at <http://dnb.d-nb.de>.

1st edition - Göttingen: Cuvillier, 2020

© CUVILLIER VERLAG, Göttingen, Germany 2020

Nonnenstieg 8, 37075 Göttingen, Germany

Telephone: +49 (0)551-54724-0

Telefax: +49 (0)551-54724-21

www.cuvillier.de

All rights reserved. This publication may not be reproduced by photomechanical means (photocopying, microfiche), in whole or in part, without the prior express permission of the publisher.

1st edition, 2020

This publication is printed on acid-free paper.

ISBN 978-3-7369-7283-4

eISBN 978-3-7369-7283-5



Contents

| | | |
|----------|---|-----------|
| 1 | Summary | 5 |
| 2 | The early history of the Δa photometric system | 7 |
| 3 | The diagnostic tools of the Δa photometric system | 8 |
| 4 | Other indices measuring the 5200Å region | 10 |
| 5 | The variability of the Δa index | 13 |
| 5.1 | CP stars | 14 |
| 5.2 | Be/shell stars - The case of Pleione | 20 |
| 5.3 | Variable star detection via Δa photometry | 21 |
| 5.3.1 | Temporal analysis and results | 22 |
| 5.3.2 | Individual variables and Conclusions | 26 |
| 6 | Chemically Peculiar stars of the upper main sequence | 27 |
| 7 | Other star groups interesting for the Δa photometric system | 30 |
| 7.1 | Blue Stragglers | 30 |
| 7.2 | Herbig Ae/Be stars | 32 |
| 7.3 | Be/shell stars | 33 |
| 7.4 | B[e] stars | 33 |
| 7.5 | Evolved and Population II stars | 35 |
| 8 | An inspection of the wavelength region from 4900 to 5700Å | 38 |
| 8.1 | The 5200Å flux depression | 38 |
| 8.2 | Features of Stars across the HRD | 40 |
| 8.3 | Features of Quasars | 48 |
| 8.4 | Features of Galaxies | 50 |
| 9 | Spectrophotometry of the 5200Å region for peculiar and normal type stars | 54 |
| 9.1 | Data and reduction | 54 |
| 9.2 | Results | 58 |
| 9.3 | Known peculiar objects | 59 |
| 9.4 | Apparent normal-type objects | 61 |
| 9.5 | Conclusions | 66 |



| | |
|---|------------|
| 10 A catalogue of Δa measurements for Galactic field stars | 67 |
| 10.1 Selection, preparation, and homogenization of the data . . . | 69 |
| 11 The detection probability of the Δa system for Population I stars | 72 |
| 11.1 The normal type objects | 73 |
| 11.2 Outliers not included in RM09 | 75 |
| 11.3 CP1 stars | 75 |
| 11.4 CP2 stars | 78 |
| 11.5 CP3 stars | 79 |
| 11.6 CP4 stars | 79 |
| 11.7 λ Bootis stars | 79 |
| 11.8 Supergiants | 79 |
| 11.9 Be/shell stars | 81 |
| 11.10 Newly confirmed CP stars | 81 |
| 11.11 Comparison with the Geneva $\Delta(V1 - G)$ and Z indices . . . | 82 |
| 12 An empirical temperature calibration for the Δa photometric system | 84 |
| 12.1 The B-type stars | 85 |
| 12.1.1 Sample of program stars | 86 |
| 12.1.2 The estimation of the reddening | 87 |
| 12.1.3 The calibration of the effective temperature | 88 |
| 12.1.4 The calibration procedure for the Δa photometric system | 89 |
| 12.2 The A-type and mid F-type stars | 92 |
| 12.2.1 Sample of program stars | 93 |
| 12.2.2 The estimation of the reddening | 94 |
| 12.2.3 The calibration of the effective temperature | 95 |
| 12.2.4 The result for the Δa photometric system | 96 |
| 13 Theoretical isochrones for the Δa photometric system | 98 |
| 13.1 Models, data selection, and isochrone fitting | 99 |
| 14 Synthetic Δa photometry | 104 |
| 14.1 The normality line | 104 |
| 14.1.1 The synthetic Δa filter system | 106 |
| 14.1.2 Calibration relations | 106 |
| 14.1.3 Luminosity effects | 110 |
| 14.1.4 Metallicity effects | 112 |
| 14.2 The cool CP and λ Bootis stars | 112 |



| | | |
|-----------|---|------------|
| 14.2.1 | Model atmospheres for CP stars | 113 |
| 14.2.2 | Selected sample of stars | 116 |
| 14.3 | Individual models: parameter determination with observed fluxes | 118 |
| 14.3.1 | CP1 stars | 119 |
| 14.3.2 | CP2 stars with $T_{\text{eff}} \leq 8500$ K | 122 |
| 14.4 | CP2 stars with $8500 \text{ K} \leq T_{\text{eff}} \leq 10000$ K | 127 |
| 14.5 | Comparison of the synthetic and observed Δa photometry for ten CP stars | 130 |
| 14.5.1 | λ Bootis stars | 130 |
| 14.5.2 | CP1 stars | 131 |
| 14.5.3 | CP2 stars | 131 |
| 14.5.4 | Spectroscopic observations of the 5200\AA depression | 133 |
| 14.6 | Observational constraints | 134 |
| 14.7 | Microturbulence, abundances, and atomic data | 136 |
| 14.8 | Stratification and magnetic fields | 137 |
| 14.9 | Stellar model atmospheres with magnetic line blanketing | 139 |
| 14.9.1 | The stellar model atmosphere code LLModels | 139 |
| 14.9.2 | The behaviour of Δa , $(\Delta V1 - G)$, and Z in correlation with T_{eff} and B | 140 |
| 14.9.3 | Effects of the individual abundance patterns | 143 |
| 15 | Heuristic observational correlations for CP stars | 146 |
| 16 | Investigating the λ Bootis spectroscopic binary hypothesis using Δa photometry | 153 |
| 16.1 | Modelling the spectroscopic binary systems | 153 |
| 16.2 | Results and Conclusions | 156 |
| 17 | The Δa observations of Galactic open clusters | 158 |
| 17.1 | The characteristics of open clusters | 158 |
| 17.2 | Photoelectric Δa observations of Galactic open clusters | 160 |
| 17.2.1 | Observations and reduction | 161 |
| 17.2.2 | Feinstein 1 | 164 |
| 17.2.3 | NGC 2168 | 165 |
| 17.2.4 | NGC 2323 | 166 |
| 17.2.5 | NGC 2437 | 166 |
| 17.3 | CCD Δa observations of Galactic open clusters | 167 |
| 17.4 | The observation of Blue Stragglers in open clusters | 170 |
| 17.5 | Statistics of the observed open clusters | 171 |



| | | |
|-----------|---|------------|
| 18 | The first Δa observations of three Galactic globular clusters | 176 |
| 18.1 | Target selection, observations and reduction | 179 |
| 18.2 | Discussion | 180 |
| 19 | The Δa observations within the Large Magellanic Cloud | 186 |
| 19.1 | Target fields and observations | 187 |
| 19.2 | Reduction and calibration | 189 |
| 19.3 | The incidence of the apparent peculiar objects | 191 |
| 19.4 | The spectroscopic verification of one CP and Be star . . . | 195 |
| 19.4.1 | Target selection, observations, and reductions . . . | 195 |
| 19.4.2 | Results and conclusions | 196 |
| 19.5 | The (non-)variability of magnetic CP candidates in the LMC | 199 |
| 19.5.1 | I light variability of mCP stars | 202 |
| 19.5.2 | Periodograms | 204 |
| 19.5.3 | Discussion of periodograms | 205 |
| 19.5.4 | The mCP candidates suspected of periodic light variations | 207 |
| 19.5.5 | Conclusions | 210 |
| 20 | A case study of using Gunn g data to improve Δa | 211 |
| 21 | The ALHAMBRA and COSMOS surveys | 215 |
| 22 | Conclusions and Outlook | 218 |
| 23 | References | 219 |



1 Summary

This book describes the history, characteristics, observations, results and future prospects of the Δa photometric system. Up to now, it successfully produced countless scientific output since its development and first appearance on the astronomical scene initiated by Hans-Michael Maitzen in 1976.

It was intended to study the typical and unique flux depression at 5200 Å found in mainly magnetic chemically peculiar stars of the upper main sequence. Therefore the most important measurement is through the g_2 filter centred at this wavelength region. In addition, one needs the information about the continuum flux of the same object. Originally, the g_1 (centred at 5000 Å) and Strömgren y (5500 Å) was used to get the continuum flux. In principle, any other filters such as Johnson $BVR I$ can be used. However, the closer it is measured to the 5200 Å region and the narrower the used filters are, the results become more accurate. The a index is then defined as the flux in g_2 compared to the continuum one. Because there is a general increase of opacity around 5200 Å with decreasing temperature, one has to normalize a with the index a_0 of a non-peculiar star of the same temperature, to compare different peculiar (or deviating) stars with each other (Δa index).

With this photometric system it is possible to measure any peculiarities, such as abnormal absorption and emission lines, in the region of 5200 Å. Soon, it was used to investigate metal-weak, Be/shell and supergiants in the Galactic field and open clusters.

At the end of the 1980ies, the highly developed photomultiplier technology, slowly but surely, was replaced by the CCD technology. It took a few years until CCDs reached the same accuracy as photomultipliers for the same exposure time and target brightness.

The first CCD Δa observations which were performed 1995 with the same telescope, the 61 cm Bochum telescope at La Silla, by the same person, Hans-Michael Maitzen, who initiated in 1971 the observations of CS Vir in g_1 and g_2 . The results of the first CCD observations, data for 25 Galactic field stars with known spectroscopic peculiarity types, were published in 1997. After that, CCD observations of open clusters, the Large Magellanic Cloud and Globular clusters were successfully secured.

This book goes one step further showing that the Δa photometric system and the study of its corresponding spectral region from 4900 to 5700 Å is able to contribute significant new insights at many fields of astrophysics. The list of objects which are potential targets is long: cool-type Population



I and Population II objects, supergiants, emission type objects, all type of galaxies, and so on.

These research fields can be investigated by 1) new observations; 2) archival data not used so far, for example cool-type objects in star cluster fields and 3) data from other surveys which employ similar filters. For sure, all three approaches will be extensively utilized in the future.



2 The early history of the Δa photometric system

The roots of the Δa photometric system date back to the year 1969. The invention and development of it is closely tied to ao. Univ.-Prof. i.R. tit. Univ.-Prof. Dr. Hans-Michael Maitzen (HMM). The interested reader is referred to his article “Two decades of Δa photometry” published by Maitzen (1998) which includes a comprehensive overview of the history until then.

Here, a very short overview is given because it sheds more light on the original intentions which are still state-of-the-art today.

In 1969, HMM got a position at the newly founded astronomical institute of the Ruhr-University of Bochum, Germany. Soon, the at that time, head of the institute, the late Prof. Theodor Schmidt-Kaler invited him to perform photometric observations at the Bochum 61 cm telescope at ESO (La Silla). He started the observations in May 1969, only less than two months after the official opening of the observatory on 23. March 1969. He monitored magnetic stars (CP2) from the catalogue by Babcock (1958) as suggested to him by the late Prof. Karl Rakos. This was to confirm the variability of these stars by means of Johnson UBV photometry. The “key star” for the further historical development was CS Vir (HD 125248), a classical CP2 object. HMM found not only, as expected, a decrease of the variability amplitude from U to B , but also the V light curve exhibited either half the period of U and B , or as we know today, a double wave variation due to the variable continuum backwarming produced by strong line absorption at shorted wavelengths.

These results were published by Maitzen & Rakos (1970). As next step, photometric observations in the Strömgren system and spectroscopy of CS Vir was done. Because no filter covered the wavelength interval where the transition from a single to a double wave variation occurred, he decided to order two new interference filters (g_1 , $\lambda_c = 5020 \text{ \AA}$ and g_2 , $\lambda_c = 5240 \text{ \AA}$, both with FWHM = 130 \AA). These filters are already very close to those used today (Sects. 9 and 14.1.1). From their colour appearance to the human eye, Prof. Schmidt-Kaler suggested to name them *giftgrün* (poisonous green, g_1) and *lindgrün* (yellowish green, g_2) which was rejected for practical reasons. The first observations with these filter were conducted in 1971 in order to trace the light variability of CS Vir but not in the way, the Δa photometric system is used nowadays. The most important result for the birth of the system was the detection of the characteristic flux depression at 5200 \AA by Kodaira (1969) which can be measured via

g_2 . This prompted HMM to substantiate the feature by a more extended photometric campaign of different CP stars (not only magnetic ones) and apparent normal type ones. Using the already available Strömgren y filter, he defined the normality line, and the final Δa parameter. The foundation paper then appeared more than 40 years ago (Maitzen, 1976a).

3 The diagnostic tools of the Δa photometric system

Here, an overview of the characteristics and diagnostic tools of the Δa photometric system is given. One has to refer to many consecutive Sections where these tools are applied and discussed in more detail. Some of the information given here might be repeated there because of the necessity to clarify some points within the context. However, it was tried to minimize such duplicities.

Basically, the Δa photometric system consists of one filter which measures the 5200Å region (g_2) and an additional information about the continuum flux of the same object. This can be either achieved by measuring the flux at the adjacent spectral regions, for example with the filters g_1 (5000Å) and $g_3 = y$ (5500Å), but also by any other effective temperature (T_{eff}) sensitive colour, $B - V$, $b - y$, or $B2 - V1$ to give some examples. The latter were indeed successfully applied, for example, Maitzen et al. (1986) used $b - y$, whereas Maitzen (1993) also employed $B - V$ colours.

With these measurements, the index a can be calculated as

$$a = g_2 - \frac{g_1 + y}{2} \quad (1)$$

$$a = g_2 - \frac{b + y}{2} \quad (2)$$

$$a = g_2 - \frac{B + V}{2} \quad (3)$$

In principle, the positioning of the continuum filters minimizes the influence of T_{eff} on the peculiarity index. The closer it is measured to the 5200Å region and the narrower the used filters are, the results become more accurate. One has to avoid, for example, the hydrogen or helium lines for the continuum measurement because they are very much sensitive to the projected rotational velocity and the surface gravity.

In Sect. 10, Table 9 lists several different filters which were used in the past for different purposes to obtain observations. There is a general increase of opacity around 5200Å with decreasing temperature (Sect. 14.1).




Figure 1: The two basis diagnostic tools of the Δa photometric system for three open clusters: the normality line a_0 versus $(g_1 - y)$ and colour-magnitude M_V versus $(g_1 - y)$ diagrams together with isochrones.

Therefore, one has to normalize a with the index a_0 of a non-peculiar star of the same temperature, to compare different peculiar (or deviating) stars with each other. The photometric peculiarity Δa index is therefore defined as

$$\Delta a = a - a_0[(g_1 - y); (b - y); (B - V)]. \quad (4)$$

The first diagnostic tool is the *normality line* which comprises of the location of the a_0 -values in respect to a T_{eff} sensitive colour. Assuming that all stars exhibit the same interstellar reddening, peculiar objects deviate from the normality line more than 3σ (Fig. 1).

From observations, it is known that the normality line is shifted by $E(b - y)$ to the red and by a small amount $E(a)$ to higher a -values (Maitzen, 1993). It has to be emphasized that correlations between the amount of reddening for different photometric systems exist (Sect. 12.1.2, Yuan et al., 2013), for example $A_V = 3.1E(B - V) = 4.3E(b - y)$. These correlations can be used to transform the reddening values between the different photometric systems. The ratio of the shifts

$$f = E(a)/E(b - y) \quad (5)$$

can be determined from the deviation of a reddened cluster normality line from the unreddened relationship. On the other hand, assuming a mean f -value (≈ 0.05) and iterating the formula

$$a(\text{corr}) = a(\text{obs}) - fE(b - y) \quad (6)$$

one can determine reddening values by the Δa -photometry of clusters. The effect has only to be taken into account for $E(b - y) > 0.3$ mag and a non uniform reddening distribution.

More of a problem is the estimation of differential reddening within a star cluster (Bonatto et al., 2012), for example. In principle, several methods can be applied (Sects. 12.1.2 and 12.2.2) to overcome this problem:

- to deredden each individual object using the Strömgen $uvby\beta$ photometric system and its calibrations
- to use the Q -method for the Johnson UBV photometric system

An a versus $(g_1 - y)_0$ or $(b - y)_0$ or $(B - V)_0$ diagram should then be able to further strengthen the membership of objects to the investigated star cluster since fore- and background deviate significantly from the normality line. These deviations are in generally ten times higher than those observed for the most prominent CP stars.

The second diagnostic tool is a y versus $(g_1 - y)$ diagram. Instead of y , any other magnitude, for example, V , R , I , and so on, can be taken. This is just a classical colour-magnitude diagram (CMD) for the Δa photometric system. With the knowledge of the reddening and the distance, the absolute magnitude M_V , and using the bolometric correction (B.C.) as well as the absolute magnitude of the Sun, the luminosity $\log L/L_\odot$ can be calculated (Gómez et al., 1998).

In Sect. 13, a grid of isochrones with different initial chemical compositions for the Δa system is presented. There, it is shown that the accuracy of fitting isochrones to Δa data without the knowledge of the cluster parameters is between 5 and 15 %. This efficient tool has been already widely used for star clusters and the LMC (Sects. 17 and 19).

Figure 1 shows M_V versus $(g_1 - y)$ diagrams (upper panels) together with isochrones for three open clusters. The capability to sort out non-members and to estimate the cluster parameters are evident.

4 Other indices measuring the 5200Å region

One has to differentiate between indices which measure the flux in the 5200Å region directly and those which measure the metallicity (line blan-




Figure 2: The filter transmission curves of the Geneva 7-colour (upper panel) and Vilnius (lower panel) system taken from Bessell (2005).

keting). The latter are capable to detect most extreme CP stars because of their prominent overabundances. Those are, for example, the m_1 , Δm_1 , and $[m_1]$ in the Strömgren system as well as m_2 and δm_2 in the Geneva 7-colour system (Golay, 1974).

Another interesting approach was published by Masana et al. (1998). They defined a reddening-depended peculiarity index Δp as a linear combination of all Strömgren $uvby\beta$ colours. They found a detection rate of up to 50% for early (hot) CP2 stars. However, this approach was not systematically followed after the original paper has been published. It would be very interesting to use the new photometric catalogue by Paunzen (2015) to update and upgrade this method.

Besides the Δa photometric system (Sect. 3), a few other indices are available which measure the flux in the 5200Å region. These are described in more detail in the following.

The $\Delta a'$ index by Adelman (1979): it basically works in the same way as the “classical Δa index”, but was especially designed for his spectrophotometric data. It is defined as

$$a' = m_{5264} - [m_{4785} + 0.453 \cdot (m_{5840} - m_{4785})]. \quad (7)$$

The measurements at 4785 and 5840Å, which were both made at locations significantly bluer than g_1 and redder than y , represent a wavelength base for the continuum that is larger by a factor of two than in Δa photometry. The depression itself is represented by the 5264Å value. Like the a index,

the a' index is slightly dependent on the colour of the star and therefore has to be normalized by the index of non-peculiar stars with the same colour.

The $\Delta(V1 - G)$ and Z indices of the Geneva 7-colour system: The Geneva 7-colour photometric system (Fig. 2) is the most homogeneous one because unique filter sets together with the same type of photomultipliers were used throughout its history (Golay, 1994). However, unfortunately, no new observations are available for it because the original instrumentation is not existing any more. The $\Delta(V1 - G)$ index is the first measurement for peculiarity derived from the Geneva 7-colour photometric system. The $V1$ and G filters are centred at 5408 and 5814Å (bandwidths of about 200Å), respectively. Hauck (1974) was the first to propose this index as peculiarity parameter. It is defined as

$$\Delta(V1 - G) = (V1 - G) - 0.289 \cdot (B2 - G) + 0.302. \quad (8)$$

On average, normal stars have $\Delta(V1 - G)$ values of -5 mmag (Sect. 11.11). The zero point of the $\Delta(V1 - G)$ index represents the upper limit of the sequence of normal type objects and not its mean value. This was done by using the upper envelope for normal type, luminosity class V to III objects, based on a linear fit for the correlation of $(V1 - G)$ with $(B2 - G)$ as given by Hauck (1974) which introduces this negative shift. Consequently, a very strict significance limit of $+10$ mmag for $\Delta(V1 - G)$ was set by Hauck & North (1982) to avoid contamination of CP objects.

Besides the $\Delta(V1 - G)$ index, Z within the Geneva 7-colour photometric system is most suitable for detecting CP stars. Originally, Cramer & Maeder (1979) defined a three-dimensional grid $[XYZ]$ for hot stars on the basis of the seven available filters (Fig. 2). The X parameter is oriented along the main sequence (MS) of O- and B-type, whereas the Y is in the direction of high luminosity stars. The Z parameters is normal to the $[XY]$ plane. It is defined (Cramer, 1999) as

$$Z = -0.4572 + 0.0255 \cdot U - 0.1740 \cdot B1 + 0.4696 \cdot B2 - 1.1205 \cdot V1 + 0.7994 \cdot G \quad (9)$$

The Z index is virtually independent of temperature and gravity effects for stars hotter than A0 or $(b - y)_0 < 0$ mag. Furthermore, the only stars showing a significant deviation in the Z direction, are the CP stars. Later on, Cramer & Maeder (1980) showed that Z is correlated with the measured magnetic field of CP stars. This is line with the corresponding synthesized photometry (Sect. 14.9.2). Cramer (1999) lists a limit of ± 10 mmag for apparent peculiarity (Sect. 11.11).

As the Δa index (Sect. 5.1), $\Delta(V1 - G)$ and Z also vary over the rotational period of CP2 stars (Muciek et al., 1984). However, the variations (± 3 mmag) are less than those for the Δa index.

The different diagrams of the Vilnius system: The Vilnius system (Fig. 2) was developed independently from the Geneva 7-colour system but for similar reasons, namely, to derive temperatures, luminosities, and peculiarities in reddening and composition from photometry alone. The colours are normalized by the condition $U - P = P - X = X - Y = Y - Z = Z - V = V - S$ for unreddened O-type stars. Therefore, all colours for normal stars are positive. Reddening free indices are constructed as for the Geneva 7-colour and Strömgen systems. Interesting enough, the Z filter was placed on the Mg I triplet as well as the MgH molecular band. It is also sensitive to the luminosity classes of G- to M-type stars (Straizys et al., 1986). Straizys & Žitkevičius (1977) proposed four diagrams to separate CP from normal type stars. They defined the following index

$$Q_{YZS} = Y - Z - 0.538 \cdot (Z - S) \quad (10)$$

which is, in principle, independent of interstellar reddening. If a star exhibits $Q_{YZS} < -0.03$ mag it can be considered as CP object. They found that 56% of CP stars (denoted as Ap stars in their work) have such values with the classical hot Si stars show the largest and the HgMn (CP3) the smallest deviations. It turned out that strongly reddened and cool type CP stars can not be unambiguously detected. Also, this index fails to detect CP1 stars which is similar to the situation within the Δa photometric system (Sect. 11.3). However, CP1 stars can be detected in the Vilnius system on the basis of CMDs including the U filter. Later on, North et al. (1982) discussed the capabilities of a joint Vilnius and Geneva 7-colour (VILGEN) system to detect CP stars. They improved the detection capability of $\Delta(V1 - G)$ by using the Vilnius Z filter (not to be confused with the Geneva Z index) instead of $V1$. In addition, they defined four peculiarity indices (PECx). Using these indices, 54% of magnetic Ap (CP2 and CP4) stars were detected. Since then, no new corresponding investigation has been published employing this extended system.

5 The variability of the Δa index

If one looks very closely on the brightness of any arbitrary chosen star, one will find variability on different time scales. The detection limit of

the amplitude only depends on the instrumentation and accuracy of the measurements. The Sun, as the closest star, shows variability with periods from minutes (not visible by eye) to several years (sunspots). Variable stars are divided into two categories: intrinsic variables, in which internal physical changes, such as pulsations or eruptions, are the driving mechanism, and extrinsic variables, in which the light output fluctuates due to planet transits, eclipses or stellar rotation (Percy, 2007). The further classification is rather complex; originally it was based on a star's light-curve characteristics, amplitude, and periodicity (or the lack of it). Many astrophysical theories, for example pulsation, diffusion, and evolutionary models, can be tested with variable stars (Handler, 2013).

It is also well known that the amplitude of almost all types of variability is depending on the observed wavelength region. For example, CP stars exhibit larger amplitudes in the UV than in the IR region (Krtička et al., 2012). Therefore, it was several times tested if and how the variability of stars influences the 5200Å region and the Δa index. The knowledge of this behaviour is essential to estimate the percentage of missed positive (or negative) detections, for example.

5.1 CP stars

Here, the published results of the analysis of the variability behaviour for 17 well established CP stars is presented. Let us recall that the main characteristics of this group are (Sect. 6): peculiar and often variable line strengths, quadrature of line variability with radial velocity changes, photometric variability with the same periodicity and coincidence of extrema. Stibbs (1950) introduced the Oblique Rotator concept of slowly rotating stars with non-coincidence of the magnetic and rotational axes. This model reproduces the photometric variability by the appearing and receding patches on the stellar surface similar to Sun spots.

Table 1 lists the spectral types (Skiff, 2016), dereddened ($b - y$) colours (Sect. 10), averaged quadratic effective magnetic field $\langle B_e \rangle$ (Bychkov et al., 2009), the rotational period P , the observed Δa range, and the mean absolute Δa value of the sample. It was chosen to list here the $\langle B_e \rangle$ values by Bychkov et al. (2009) because they compiled a homogeneous sample of measurements. For the definition of $\langle B_e \rangle$ and its correlation with the magnetic field modulus, for example, the reader is referred to Bychkov et al. (2003). Briefly, the effective magnetic field B_e of a star is a complex average over the stellar disc of the projection of the local vector of magnetic intensity on the line of sight, and can be measured as the splitting of circularly polarized components of spectral lines.

Table 1: The spectral types (Skiff, 2016), dereddened $(b - y)$ colours (Sect. 10), averaged quadratic effective magnetic field $\langle B_e \rangle$ (Bychkov et al., 2009), the rotational period P (references see text), the observed Δa range (references see text), and the mean absolute Δa value (Sect. 10). For some objects, the mean absolute Δa value could be higher than the given range because there is an offset between different applied filter systems (see Section 7 in Hensberge et al., 1981).

| HD | HIP | Spec | $(b - y)_0$ [mag] | $\langle B_e \rangle$ [G] | P [d] | Δa_{range} [mmag] | Δa [mmag] |
|--------|-------|--------------|----------------------|------------------------------|------------|-------------------------------------|----------------------|
| 3980 | 3277 | Ap SrEuCr | +0.079 | 780(25) | 3.9516 | +39 ... +49 | +38 |
| 5601 | 4488 | Ap Si | -0.056 | 1192(100) | 1.11 | +52 ... +66 | +49 |
| 19712 | 14736 | A2p CrEu | -0.060 | 2268(225) | 2.1945 | +44 ... +62 | +43 |
| 25267 | 18673 | B9p Si | -0.070 | 241(91) | 1.21 | +30 ... +50 | +36 |
| 30466 | 22402 | B9p SiCr | +0.025 | 1716(354) | 2.7795 | +42 ... +57 | +50 |
| 30849 | 22340 | Ap SrCrEu | +0.166 | | 15.865 | +12 ... +37 | +30 |
| 38823 | 27423 | A5p Sr | +0.151 | 1509(111) | 8.635 | +06 ... +16 | +19 |
| 50169 | 32965 | Ap SrCrEu | -0.035 | 1218(70) | 1.729 | +65 ... +67 | +78 |
| 52847 | | Ap CrEuSr | +0.102 | | | +53 ... +58 | +62 |
| 53116 | 34049 | Ap SrEu | -0.046 | | 11.978 | +38 ... +63 | +50 |
| 55540 | | Ap CrEu | -0.068 | | | +66 ... +71 | +70 |
| 56022 | 34899 | Ap SiSr | -0.022 | 202(117) | 0.9184 | +18 ... +26 | +16 |
| 72968 | 42146 | A0p SrCr | -0.039 | 445(181) | 11.305 | +12 ... +21 | +50 |
| 81009 | 45999 | A2p SrCrEuSi | +0.086 | 1431(204) | 33.984 | +29 ... +37 | +35 |
| 111133 | 62376 | A0p SrEuCr | -0.057 | 807(143) | 16.31 | +43 ... +73 | +56 |
| 116458 | 65522 | Ap CrSrEu | -0.036 | 1926(273) | 126.18 | +14 ... +32 | +54 |
| 126515 | 70553 | A2Vp SrSiCr | -0.038 | 1859(360) | 129.95 | +33 ... +74 | +52 |

Note that the $g_1, g_2, g_3/y$ filters, for example, used by Hensberge et al. (1981) are slightly different than those used by Maitzen (1976a). Therefore, there is an offset between these systems (see Section 7 in Hensberge et al., 1981). In Sect. 10, mean absolute Δa values from different sources are presented. These values can be, for some objects, higher than those derived by the references mentioned in the following. However, the aim of this analysis is, if the individual CP stars would have been detected independent of the knowledge about the observed phase of variability. This is the most general case possible. Or in other words, is the expected minimum Δa value over the phase still above the detection limit (Sect. 10). In the following, the results of the individual stars, listed in Table 1, are discussed in more detail.

HD 3980: Nesvacil et al. (2012) performed a spectrum analysis in order to determine atmospheric parameters for Doppler imaging. They detected also horizontal inhomogeneities (stratification) in the stellar atmosphere. However, no obvious correlation between theoretical predictions of diffusion

in CP stars and the abundance patterns could be found. The rotational period of 3.9516 d was deduced by Maitzen et al. (1980). They also investigated the behaviour of the Δa index over the rotational period (+39 ... +49 mmag).

HD 5601: Besides the classification as CP star, not many investigations were devoted to it. Joshi et al. (2009) unsuccessfully search for rapid oscillations (periods below 25 minutes) whereas Hensberge et al. (1981) estimated a rotational period of close to one day. The Δa index varied for ± 7 mmag on a very high absolute level.

HD 19712: The rotational periods published by Hensberge et al. (1981, 2.1945(2) d) and Dubath et al. (2011, 2.0422 d) are slightly different. The latter is based on the Hipparcos photometry (Høg et al., 2000). However, Hensberge et al. (1981) noticed that one of their apparent constant comparison star, HD 20319, seems to be a low amplitude variable. Speckle interferometry (Horch et al., 2006) revealed that HD 20319 is a close binary system with a separation of about 1". The variability of this object could be due to binarity. For HD 19712, the variations of Δa range from +44 to +62 mmag.

HD 25267: Glagolevskij & Nazarenko (2015) presented a detailed analysis of the magnetic field geometry for this star with its period slightly longer than one day. The rather large variability of the Δa index (Hensberge et al., 1981) can be possibly explained by the strong dipole shift across the axis when the magnetic poles are located close to each other (Glagolevskij & Nazarenko, 2015).

HD 30466: It is a double wave photometric variable CP2 star ($P = 2.7795$ d) with an averaged quadratic effective magnetic field of about 1.5 kG (Bychkov et al., 2009). The measured Δa values over the rotational period are between +42 and +57 mmag (Maitzen, 1976b) which are well above detection limit.

HD 30849: This star was often used as a “standard object” which does not show any rapid oscillation (Balmforth et al., 2001). It is one of the coolest CP star investigated for variations in the Δa index. The rotational period of about 15.865 d (Hensberge et al., 1982) is well established. The lower measured Δa of +12 mmag (Hensberge et al., 1981, 1982) is close to the defined detection limit.

HD 38823: Hensberge et al. (1981) found a rotational period of 8.635 d for this cool CP star with a rather strong averaged quadratic effective magnetic field (1.5 kG, Bychkov et al., 2009). The maximum observed Δa index of +19 mmag (Sect. 10) is not very high, probably because of its rather low T_{eff} and the decreasing sensitivity of Δa in this regime (Sect. 14.9.2). In certain phases of the rotational cycle, HD 38823 would have

been not detected photometrically.

HD 50169: There is variability of a very long time scale (at least 1200 d) and one which should be connected to the rotational period (1.729 d) present (Heck et al., 1987). However, none of these periods is firmly settled. Adelman et al. (1998), on the other hand, found no variations in their own Strömgren *uvby* as well as Hipparcos photometry. Therefore, the characteristics of the variability still remains unsolved. But due the well established CP classification (Adelman, 1981) and the measured magnetic field, one should expect stellar spots and rotational induced variability. This conclusion is support be the very large Δa value of +78 mmag (Sect. 10).

HD 52847: Up to now, no evidence for variability of this star was found (Maitzen, 1976a; Hensberge et al., 1981). The measured range of Δa values was interpreted as intrinsic scatter. Maitzen (1976a) deduced $\Delta a = +63(4)$ mmag from 16 measurements whereas Hensberge et al. (1981) presented only three measurements. Freyhammer et al. (2008) confirmed the CP nature of HD 52847 and also found a strong mean magnetic field modulus of 4.4 kG. Based on photometric data from the All Sky Automatic Survey (ASAS, Pojmanski, 2002) they established an upper limit for the amplitude of variability of 4.3 mmag.

HD 53116: Hensberge et al. (1981, 1982) found a rotational period of 11.978 d with evidence for an antiphase relation of variations for Strömgren *v* with *uby* (Vogt & Faúndez, 1979). The presence of a strong magnetic field was only concluded on the basis of photometric data (North & Kroll, 1989). The significant variations not only in Δa , but also within Strömgren *uvby* strongly supports that a strong magnetic field is present causing large surface spots.

HD 55540: Freyhammer et al. (2008) detected a strong mean magnetic field modulus of 12.7 kG for this object. There are no other investigations of this star available in the literature which shows that many members of this group are still hardly investigated. Maitzen (1976a) list $\Delta a = +70(5)$ mmag and found no signs of variability. Hensberge et al. (1981) reported significant differences (40 mmag) in Strömgren *ub* on a long time base. No other investigations in this respect are available in the literature.

HD 56022: This is a CP2 star with a rather weak averaged quadratic effective magnetic field and a well established period of 0.9184 d. Hempelmann & Schöneich (1988) presented a detailed analysis of its light curve in the UV region including the spot characteristics using data from the TD-1 (Humphries et al., 1976) and ANS (Wesselius et al., 1982) satellite experiments. HD 56022 is one of the rare CP2 stars for which the “null wavelength” λ_0 , defined where for $\lambda < \lambda_0$ a light curve is in counterphase to

a curve $\lambda > \lambda_0$, is between 2500 and 3300 Å (NUV region). Hensberge et al. (1981) found no significant variations in Strömgren *vby* but for *u*. This is in line with the results by Hempelmann & Schöneich (1988) who found the largest amplitudes in the Far-UV with a decrease to the Near-UV. The absolute Δa value (+16 mmag) and its variations are very small and can be explained by the above mentioned characteristics.

HD 72968: Maitzen et al. (1978) presented extensively photometric data for this double wave variable CP2 star with a period of 11.305 d. The listed observations (see Table 1 therein) were used to calculate Δa over the period, resulting in a range between +12 and +21 mmag. This low values can be explained by the rather weak averaged quadratic effective magnetic field (500 G, Bychkov et al., 2009) of this star. According to the detection limit deduced in Sect. 11, it would be recognised as CP star candidate only in certain phases.

HD 81009: This is a very close visual binary system with a separation of the components ($\Delta V = 1.2$ mag) between 0.1 and 0.2" (Tokovinin et al., 2015). The properties of this very interesting system was investigated in more detail by Wade et al. (2000). They found that HD 81009 is a highly eccentric ($\epsilon = 0.718$), long-period ($P_{\text{orb}} = 29.3$ yr) binary system composed of two MS A-type stars. The hotter primary component was identified as the slowly rotating ($P_{\text{rot}} = 33.984$ d) magnetic CP2 star. The longitudinal magnetic field strength varies between 6.8 and 9.8 kG. The photometric variations presented by Hensberge et al. (1981) have the highest amplitude in Strömgren *v*. No significant light variations were detected in Strömgren *y* and Δa .

HD 111133: Again, a CP2 star with a moderate strong averaged quadratic effective magnetic field of about 1 kG (Bychkov et al., 2009) and a rotational period of 16.31 d (Buchholz & Maitzen, 1979). The latter investigated this object in the region between 4800 and 5600 Å using a spectrum scanner. They observed Δa ten times over the complete phase. The values ranged between +43 and +73 mmag which makes this star to one with the largest measured Δa values.

HD 116458: This is a spectroscopic binary system containing at least one CP star with an orbital period of 126.18 d (Ren & Fu, 2013). There was also a 2 kG strong averaged quadratic effective magnetic field detected (Bychkov et al., 2009). It is a very unusual and peculiar object showing strong variations of Co, Hg, Mn, and Si (Dworetzky et al., 1980). The original mean Δa measurement was +54 mmag (Maitzen, 1976a). Measurements at two different phases by Maitzen & Wood (1977) resulted in a relative Δa value variation between 14 and 32 mmag, respectively.

HD 126515: Glagolevskij (2005) modelled the magnetic field variations

Table 2: The observed magnitude ranges, defined as the difference between the lowest and highest value over the rotational period, for Δa , Strömgren *uvby*, and Johnson *V*, if available. For HD 111133 and HD 126515, no additional data to the Δa ones are available.

| HD | Δa [mmag] | <i>u</i> [mmag] | <i>v</i> [mmag] | <i>b</i> [mmag] | <i>y</i> [mmag] | <i>V</i> [mmag] |
|--------|----------------------|--------------------|--------------------|--------------------|--------------------|--------------------|
| 3980 | 10 | 39 | 85 | 56 | 21 | |
| 5601 | 14 | 54 | 43 | | | 22 |
| 19712 | 18 | 44 | 34 | 30 | 19 | 41 |
| 25267 | 20 | 76 | 60 | 29 | 24 | 51 |
| 30466 | 15 | | 61 | 55 | 31 | 28 |
| 30849 | 25 | 51 | 92 | | | 20 |
| 38823 | 10 | 70 | 130 | | | 24 |
| 50169 | 2 | 5 | 6 | 5 | 7 | 4 |
| 52847 | 5 | 11 | 11 | 24 | 6 | 8 |
| 53116 | 25 | 107 | 69 | 58 | 103 | 56 |
| 55540 | 5 | 45 | 9 | 4 | 10 | 17 |
| 56022 | 8 | | | | 19 | |
| 72968 | 9 | 34 | 25 | 33 | 35 | |
| 81009 | 8 | 17 | 56 | 18 | 7 | |
| 116458 | 18 | 86 | 67 | 63 | 54 | |

($P = 129.95$ d) assuming a strongly decentred magnetic dipole in order to achieve the best agreement between the calculated and observed phase relations. He found that the field structure is determined by a barlike dipole with a considerable distance between the monopoles. Hensberge et al. (1986) presented a detailed investigation of the Δa variability in correlation with the different magnetic field components. Their most important conclusion was that magnetic intensification effects do not control the Δa strength.

Conclusions: Table 2 compares the observed magnitude ranges, defined as the difference between the lowest and highest value over the rotational period, for Δa , Strömgren *uvby*, and Johnson *V*, respectively. In general, it can be concluded that absolute differences for Δa are much smaller than those for the other filters. This can be easily understood by the definition of Δa as measuring the flux at 5200\AA and at the adjacent spectral regions (Sect. 3). Therefore, it is the difference of the amplitudes of the light variations at 5200\AA and the mean of the adjacent regions. The effect of

rotation is therefore minimized.

The total sample of investigated stars includes 17 CP2 objects (Table 1). This means that one star equals to 5.9% of the total sample. Since CP2 stars are normally slowly rotating in comparison to their apparent normal type counterparts (North, 1984), the rotational periods (1 to 130 d) of the sample represent the whole group very well. Also the magnetic field strengths seem to cover the range of the majority of CP2 stars (Bychkov et al., 2009). If one assumes the detection limit of +15 mmag which can be even easily achieved for much fainter Galactic open cluster members (Sect. 17.2.1), four stars (HD 30849, HD 38823, HD 72968, and HD 116458) would have been missed at certain phases of their rotational periods. The lower limits for these objects are +12, +6, +12, and +14 mmag, respectively, which is very close, but still below, the detection limit. One also have to emphasize that most targets with already very high absolute Δa values were chosen in order to study the possible effects of rotation.

The mean variation for all 17 stars is $\pm 8(5)$ mmag with a median value of ± 7 mmag, respectively. Taking these values and looking at the absolute values in Table 1, one can assume that the probability of a non-detection of an individual Galactic CP star due to a single observation at the minimum phase is about 10%. This probability can be significantly lowered by conduction individual observations during several nights and taking averaging values. The situation in the LMC on the basis of the available photometry is different (Sect. 19.5). There it was concluded that the rotationally modulated variability of the studied mCP candidates in the V and I bands is very weak (if present at all). It is very likely that the true amplitudes are much lower than 10 mmag, which is the limit derived from the time series analysis of the brightest mCP candidate. If one keeps in mind that the amplitude in Δa is much less than in the V and I bands, the probability of a non-detection is even significantly lower than in the Milky Way.

5.2 Be/shell stars - The case of Pleione

Maitzen & Pavlovski (1987a) presented Δa measurements of the Be/shell star HR 1180 (Pleione) within the Pleiades. It is a B8Vpe star known to exhibit prominent long-term photometric and spectroscopic variations with a period of about 35 yr (Tanaka et al., 2007). In the recent 70 years, Pleione showed twice a cyclic change between a Be-shell phase and a Be phase. For example, it entered a Be-shell phase in 1972, initiated by the appearance of a wide and shallow absorption of the Ca II K line. Then, the star developed many shell absorption lines in its spectrum. At the same time, the star had shown a decrease in its brightness. Pleione is known to